

Locus	Primer name	Primer sequence	Length (bp)	PCR cycling	Reference
<i>atpA</i>	1F*	CTDAATTCHACNGAAATYAGY	1322	5 min 95°C 3 x (45s 95°C, 2 min 55°C, 1 min 72°C) 30x (20 s 95°C, 1 min 55°C, 1 min 72°C) 7 min 72°C.	Thompson <i>et al.</i> , 2007
	2F	GCNATGGGBGAATAYTTCG			
	3R	CGGAARTATTCVCCCATNGC			
	4R*	TTACCARGWYTGGGTTGC			
	5R	GCHAGHGCDGTACGRATACC			
	6F	AGCGAWCTRATYAARCARCG			
<i>fstZ</i>	75F	GCTGTTAACACATGGTACG	750	5 min 94 °C, 30 x	Sawabe <i>et al.</i> , 2007
	800R	GCACCAGCAAGATCGATATC		(1 min 94 °C, 1min 50.5 °C, 1 min 72 °C), 7 min 72 °C	
<i>gapA</i>	150F	AACTCACGGTCGTTCAAC	750	5 min 94°C	Sawabe <i>et al.</i> , 2007
	899R	CGTTGTCGTACCAAGATAC		30x (1 min 94°C, 1 min 55°C, 1 min 72°C) 10 min 72°C.	
Y-HSP60	YHSP60-1	GACGTNGTAGAAGGTATGYAG	635	5 min 94 °C, 35 x	Kotetishvili <i>et al.</i> , 2005
	YHSP60-2	CGCCGCCAGCCAGTTAGC		(45 s 94 °C, 45 s 48 °C, 1 min 72 °C), 5 min 72 °C	
<i>pyrH</i>	4F	ATGASNACBAAYCCWAAACC	599	5 min 95°C 3 x (1 min 95°C, 2 min 15 s 50°C, 1 min 15 s 72°C) 30x (35 s 95°C, 1 min 45 s 50°C, 1 min 15 s 72°C) 7 min 72°C.	Thompson <i>et al.</i> , 2005
	2R	GTRAABGCNGMYARRTCCA			
<i>rctB</i>	rctBs	ATHGARTTYACNGAYTTYCARYNCAY	610	5 min 95°C 3 x (1 min 95°C, 2 min 15 s 55°C, 1 min 15 s 72°C) 30x (35 s 95°C, 1 min 45 s 55°C, 1 min 15 s 72°C) 7	Pascual <i>et al.</i> , 2010
	rctBas	YTTNCTYTGHATNGGYTCRAAYTCNCC			
	RTC				

				min 72°C.	
recA	1F*	TGARAARCARTTYGGTAAAGG	783	5 min 95°C 3 x (45s 95°C, 2 min 55°C, 1 min 72°C) 30x (20 s 95°C, 1 min 55°C, 1 min 72°C) 7 min 72°C.	Thompson <i>et al.</i> , 2004
	2R*	RTCRCCTTTRTAGCTRTACC			
	3F	FTYGGGBTGATGTTYGGTAACC			
	4R	GGGTTACCAACATCACVCC			
rpoA	1F*	ATGCAGGGTTCTGTDACAG	931	5 min 95°C 30x (1 min 95°C, 2 min 15 s 55°C, 1 min 15s 72°C) 7 min 72°C.	Thompson <i>et al.</i> , 2005
	3R*	GHGCCARTTTCHARRCGC			
	5F	GCAGCDCGTGTWGARCARCG			
	6R	RCGYTGYTCWACACGHGCTGC			
rpoD	70F	ACGACTGACCCGGTACGCATGTAYATG MGNGARATGGGNACNGT	780	5 min 95°C, 30x (1 min 94°C, 45s 59°C, 2 min 72°C) 10 min 72°C	Pascual <i>et al.</i> , 2010
	70R	ATAGAAATAACCAGACGTAAGTTNGC YTCNACCATYTCYTTYTT			
	400F 1200R	GAGATCATCGGTGGTGATG GAAGGACGAATCGCTTCGTG			
			800	5 min 95°C, 30x (1 min 95°C, 45s 50°C, 1 min 72°C) 10 min 72°C.	Sawabe <i>et al.</i> , 2007

(*): Indicates that those are the primers used for amplifying the gene. The other primers corresponding to this particular gene were internal, used to sequence the gene.

Kotetishvili M, Kreger A, Wauters G, Morris J, Sulakvelidze A and Stine C (2005) Multilocus Sequence typing for studying genetic relationships among *Yersinia* species. *J Clin Microbiol* 43:2674-2684.

Pascual J, Macián MC, Arahal DR, Garay E and Pujalte MJ (2010) Multilocus sequence analysis of the central clade of the genus Vibrio by using the 16S rRNA, recA, pyrH, rpoD, gyrB, rctB and toxR genes. *Int J Syst Evol Microbiol*. 60:154-65.

Thompson CC, Thompson FL, Vandemeulebroecke K, Hoste B, Dawyndt, P and Swings J (2004) Use of recA as an alternative phylogenetic marker in the family *Vibrionaceae*. *Int J Syst Evol Microbiol* 54: 919-929.

Thompson FL, Gevers D, Thompson CC, Dawyndt P, Naser S, Hoste B, Munn CB and Swings J (2005). Phylogeny and molecular identification of Vibrios on the basis of Multilocus Sequence Analysis. *Appl Environ Microb* 71: 5107-5115.

Thompson CC, Thompson F L, Vicente A C and Swings J (2007). Phylogenetic analysis of vibrios and related species by means of atpA gene sequences. *Int J Syst Evol Microbiol* 57: 2480-2484.

Sawabe T, Kita-Tsukamoto K and Thompson FL (2007) Inferring the evolutionary history of vibrios by means of Multilocus Sequence Analysis. *J Bacteriol* 21: 7932-7936.